

High Quality X-Ray and Gamma-Ray Sources

C H KEITEL¹, J EVERS¹, K Z HATSAGORTSYAN¹, AND M TAMBURINI¹

¹*Theory Division, Max Planck Institute for Nuclear Physics, Saupfercheckweg 1, Heidelberg, Germany*
Contact Email: keitel@mpi-hd.mpg.de

Recent progress is reported for the generation of high-frequency light with favorable features like narrow bandwidth, flux, coherence and polarization. This includes nuclear interaction with moving targets [1], beam-multifoil collisions [2], beam-laser collisions [3] and beam-crystal interaction [4].

References

- [1] K P Heeg, A Kaldun, C Strohm, P Reiser, C R Ott, R Subramanian, D Lentrodt, J Haber, H-C Wille, S Goerttler, R Ruffer, C H Keitel, R Röhlsberger, T Pfeifer, and J Evers, *Science* **357**, 375 (2017)
- [2] A Sampath, X Davoine, S Corde, L Gremillet, M Gilljohann, M Sangal, C H Keitel, R Ariniello, J Cary, H Ekerfelt, C Emma, F Fiuza, H Fujii, M Hogan, C Joshi, A Knetsch, O Kononenko, V Lee, M Litos, K Marsh, Z Nie, B O'Shea, J R Peterson, P S M Claveria, D Storey, Y Wu, X Xu, C Zhang and M Tamburini, *Phys. Rev. Lett.* **126**, 064801 (2021)
- [3] Y Li, R Shaisultanov, Y Chen, F Wan, K Z Hatsagortsyan, C H Keitel and J Li, *Phys. Rev. Lett.* **124**, 014801 (2020)
- [4] O D Skoromnik, I D Feranchuk, J Evers and C H Keitel, arXiv:190908046 (2019)